

TECHNICAL REPORT

70-48-FL

AD

**STORAGE LIFE OF LETTUCE AS AFFECTED
BY CONTROLLED ATMOSPHERE SYSTEM**

by

Abdul R. Rahman

Glenn Schafer

George R. Taylor

and

Donald E. Westcott

January 1970

**UNITED STATES ARMY
NATICK LABORATORIES
Natick, Massachusetts 01760**



Food Laboratory

FL 106

This document has been approved for public release and sale;
its distribution is unlimited.

Citation of trade names in this report does not constitute
an official indorsement or approval of the use of such items.

Destroy this report when no longer needed. Do not return it
to the originator.

This document has been approved
for public release and sale; its
distribution is unlimited

AD _____

TECHNICAL REPORT
70-48-FL

STORAGE LIFE OF LETTUCE AS AFFECTED BY
CONTROLLED ATMOSPHERE SYSTEM

by

Abdul R. Rahman
Glenn Schafer
George R. Taylor
Donald E. Westcott

January 1970

Project reference
2270.3

Series: FL 106

Food Laboratory
U.S. ARMY NATICK LABORATORIES
Natick, Massachusetts 01760

Foreword

Spoilage has been reported in lettuce shipped to Military installations overseas. Causes for the spoilage included slime, decay, mold, and discoloration. An investigation has been conducted on the effect of Oxytrol* modified atmosphere system, used for the transportation of fresh lettuce in refrigerated containers, on shelf life and overall quality.

This work was performed under Production Engineering Project, 2270.3.

The authors wish to acknowledge the assistance of Dr. Thomas White and the Oxytrol Corporation, Burlingame, California, for providing the Oxytrol containers, the nitrogen gas and one lettuce shipment from California to Boston. Acknowledgement is also due to Mr. O Stark for his assistance in the gas analysis and L. Dame for his monitoring of the equipment.

"Tradenames and manufacturing names are used in this report for identification purposes only and such use does not constitute indorsement or approval of any particular product, process or manufacture."

* Oxytrol is a registered Trademark of Occidental Petroleum Corporation.

TABLE OF CONTENTS

Foreword	ii
List of Tables	iv
List of Figures	iv
Abstract	v
Introduction	1
Review of Literature	1
Experimental Procedures	2
Results and Discussion	4
References	6

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	Front View - Oxytrol Model 050 Portable Container Operating Controls	8
2	Technological Evaluation Sheet	9
3	Effect of Storage Conditions on the Quality of Lettuce	10

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Effect of Storage Conditions on the Quality of Lettuce Stored 5 to 7 Weeks After Picking	11
2	Effect of Storage Conditions on the Quality of Lettuce Stored 21 to 39 Days After Purchase	12
3	Effect of Storage Conditions on the Quality of Lettuce Stored for 14 to 34 Days After Purchase	13
4	Effect of Storage Conditions on the Quality of Lettuce Stored for 2 to 6 Weeks After Purchase	14

ABSTRACT

The effect of Oxytrol controlled atmosphere system on the shelf life of lettuce was evaluated. Lettuce stored under Oxytrol at oxygen levels ranging from 3 to 5.8 percent and temperatures from 34 to 36°F for 2 to 7 weeks gave significantly higher edible yield and showed higher mean scores for overall quality than lettuce stored under normal atmosphere at similar temperatures.

INTRODUCTION

The procurement and supply of fresh produce to the Military Services overseas represents a sizable investment particularly when one considers not only the acquisition cost but also the cost of inspection, transportation, cold storage and distribution to the customer. The supply of high quality produce to the overseas customer is essential both for morale purposes and for good nutrition. However, reports continue to be received of spoilage of produce, particularly lettuce, at overseas supply points. Spoilage of lettuce is a problem which has existed for many years. Although significant improvements have been made in packaging, refrigeration, and controlled atmosphere shipping. Spoilage losses are still encountered in lettuce shipped overseas.

REVIEW OF LITERATURE

A significant amount of research has been conducted in an attempt to prolong the shelf life of fresh produce. The bulk of this research was directed toward civilian market requirements of 1-2 weeks. However, recent work was conducted involving longer storage periods ranging from 4 to 8 weeks to meet the military requirements for shipping lettuce and other fresh produce overseas.

Effects of refrigeration temperature, packaging, chemical additives, and controlled atmosphere on the shelf life of lettuce have been reported in the literature. The deterioration of lettuce is affected primarily by temperature, relative humidity and spoilage organisms. However, russet spotting and pink rib, which appear to be nonpathogenic, frequently cause severe losses even when lettuce is held under desirable transit or storage conditions (Lipton, 1961). When trimmed lettuce was stored in lined crates at 35°F. for 6 weeks, 81 percent remained edible in contrast to 61 percent of untrimmed lettuce stored in unlined crates which was edible. Lettuce retained good quality for considerably longer periods at 32°F. than at 38°F. or 45°F., regardless of the packaging materials. However, packaging with polyethylene bags was the most effective at maintaining good quality (Parsons, 1959, 1965).

Recommendations have been made that lettuce should be held at temperatures as near 32°F. as possible during the entire marketing period, since the rate of deterioration increases rapidly as temperatures rise. It is about five times greater at 75°F. than at 32°F. (Lipton 1965 and Pratt 1954). Stewart and Harvey (1967) found that the quality of lettuce at the time of unloading varied inversely with the temperature at which the lettuce had been maintained during the test. Lettuce maintained at 35°F. to 37°F. was rated "excellent" but that maintained at 38°F. or 39°F. was rated only "good". Bratley (1945) reported a disorder described as "a browning of interior leaves" observed in western grown lettuce. This may have been russet spotting, since after 6 weeks storage at 34°F. russet spotting likely would be present. Rood (1956) indicated that russet spotting can be induced artificially by exposing head lettuce to ethylene.

Results of research on the effect of controlled atmosphere on the quality of lettuce vary. Controlled atmosphere tests with lettuce showed no effects from storage in oxygen concentrations as low as 1 percent for up to 8 days at 41°F. High concentrations of carbon dioxide, however, were harmful to lettuce (Watada, 1964). Lettuce stored at 33°F., and held at 1 or 0 percent oxygen developed less russet spotting than lettuce held in air (Parsons, 1964). Littlefield et. al. (1966) reported that controlled atmospheres, within the limits of concentrations usable for fruit storage (2 percent oxygen and 2-11.5 percent CO₂), are effective fungestatic agents.

Some definite, but limited, benefits have been derived from the use of certain low O₂ atmospheres under conditions simulating the transcontinental shipment of head lettuce. The substantial reduction in russet spotting, even at O₂ concentrations as high as 8 percent, is clearly the major benefit. However, O₂ levels at 0.5 percent or lower during storage within the temperature range studied (36°F. to 50°F.), sometimes injured the heart leaves (Lipton, 1967).

Lugg (1969) stated that the use of carbon monoxide (.25 to .50 percent), in conjunction with low oxygen levels, would significantly retard butt end discoloration of lettuce. However, the expense of maintaining this level of CO with on-board tanks was expensive. But when modified atmospheres are used correctly on fruits and vegetables it was economically advantageous by providing a longer shelf life. White (1968) indicated that low oxygen levels significantly reduced decay and improved overall appearance of lettuce after 28 days of storage. Low O₂ atmospheres (3.3-3.8 percent) in liquid nitrogen refrigerated piggyback trailers significantly reduced russet spotting. Decay and butt discoloration were slightly less severe in lettuce from the nitrogen refrigerated trailer than that from the mechanically refrigerated trailers (Steward, 1968).

El-Mansy (1967) indicated that the post harvest treatment of lettuce with 6-furfurylaminopurine extended shelf life beyond that of the untreated. Gorfien et al. (1969) stated that when controlled atmospheres were maintained, significant improvements in quality and storage life were obtained. Reduction in slime, pink rib and russet spotting were found.

EXPERIMENTAL PROCEDURES

Four different experiments using different crops of Iceberg lettuce were conducted during the course of this study. The lettuce was stored in containers using the Oxytrol system. This system is a complete, self-contained atmosphere control system designed to be used as an adjunct to normal refrigeration equipment in conventional transport vehicles. These containers are designed for shipment or fixed storage of perishable commodities under low oxygen atmospheres. Liquid nitrogen is used to reduce the oxygen level. The container is insulated and equipped with a refrigeration unit to control temperature. A supply of liquid nitrogen is carried in a portable Linde LS-160B container (Figure 1). Automatic operation of the N₂ flow is provided by the special controls in the system.

The lettuce was tested at intervals ranging from 3 to 10 days by three experienced judges using the following procedure:

- a. Overall appearance - using a 9-point scale (9 = Field fresh; 1 = Inedible) (Figure 2).
- b. Defects, such as decay, pink rib, russet spotting, using a 9-point scale (9 = defects absent; 1 = severe) (Figure 2).
- c. Edible yield, determined by removing the butts as well as the defective parts of the lettuce head and weighing the remainder as the edible portion. This was divided by the original weight to obtain percent edible yield.

Oxygen levels were monitored during the experiments. A representative sample of the atmosphere in the controlled atmosphere container was analyzed chromatographically weekly. After the sample was taken, the container door was opened for few minutes to check the voltage of the oxygen analyser controller.

Experiment No. 1 - Lettuce (16 boxes; 24 heads per box) was obtained the day of harvest at Salinas, California, vacuum cooled, and shipped to the U.S. Army Natick Laboratories in two refrigerated trailers. One was equipped with Oxytrol controlled atmosphere units set at 4% oxygen. Half the lettuce in each trailer was trimmed, wrapped in polystyrene film and packed in perforated fiberboard boxes. The remainder of the lettuce was untrimmed, unwrapped and packed in slotted fiberboard boxes. The temperature of the trailers was set at $35 \pm 2^{\circ}\text{F}$. The pulp temperature of the lettuce upon arrival was 38°F . Shipping time was seven days. The 16 boxes of lettuce represented the following variables:

- a. Trimmed, wrapped lettuce shipped via Oxytrol system.
- b. Untrimmed, unwrapped lettuce shipped via Oxytrol system.
- c. Trimmed, wrapped lettuce shipped via refrigerated trailer.
- d. Untrimmed, unwrapped lettuce shipped via refrigerated trailer.

Upon receipt, the lettuce was stored in two containers 72" long, 32" wide and 89" high. The 8 boxes shipped in Oxytrol trailers were placed in an Oxytrol test container and 8 boxes shipped in normal atmosphere trailers were placed in a normal atmosphere test container. The temperature for both containers was set at 34°F . and the oxygen level in the Oxytrol controlled atmosphere container was set at 5 percent. The duration of this experiment was seven weeks, the first five in Oxytrol containers with controlled atmosphere and then two weeks in walk-in chill box at 40°F . under normal atmosphere.

Experiment No. 2 - Sixteen fiberboard cartons of naked pack (untrimmed, unwrapped) lettuce, grown in Arizona, was procured locally. The lettuce had been shipped in refrigerated containers. The lettuce was placed in a walk-in 40°F. chill box for 17 days, then placed in the Oxytrol containers at 34°F. with the oxygen level at 5 percent for 22 days.

Experiment No. 3 - Sixteen boxes of naked pack (untrimmed, unwrapped) lettuce grown in California and shipped in refrigerated trailers were purchased locally. The lettuce was placed in Oxytrol containers at 34°F. with oxygen level at 5 percent for 14 days. It was then transferred to a walk-in 40°F. chill box for 22 days.

Experiment No. 4 - Naked pack (untrimmed, unwrapped) lettuce grown in Arizona and shipped in refrigerated trailers was purchased locally. The lettuce was placed in the Oxytrol containers at 34°F. with oxygen level at 5 percent. Storage was for six weeks. The doors of the containers were opened for a few minutes at the end of each of the first two weeks to check the voltage of the oxygen analyser controller. During the last four weeks, the doors were not opened at all to determine whether the Oxygen level would drop to below 1 percent. A perforated paper bag containing two pounds of quick lime was placed in each container in order to reduce the CO₂ content.

RESULTS AND DISCUSSION

Results of experiments conducted on the storage of 4 different lettuce crops for periods from 2 to 7 weeks are shown in Tables 1, 2, 3 and 4. They indicate that lettuce stored under Oxytrol C.A. (Controlled atmosphere) where O₂ level ranged from 3 to 5.8 percent and temperatures ranged from 34°-36°F.* showed significantly higher edible yield than the ones stored under normal atmosphere at similar temperatures. Mean scores for general appearance and defects in terms of slime, mold, russet spotting, pink rib and other discolorations were higher for lettuce stored under Oxytrol controlled atmosphere system than the ones stored under normal atmosphere.

The results of experiment No. 1 are shown in Table 1. The wrapped and unwrapped field fresh lettuce stored 6 weeks after picking under normal atmosphere was a total loss, whereas 28.8 percent edible yield for unwrapped and 60.6 percent for wrapped lettuce were obtained under Oxytrol C.A. Furthermore, upon extending the storage period to 7 weeks after picking, 16.0 and 21.4 percent edible yield were obtained for unwrapped and wrapped lettuce, respectively. In general, wrapped lettuce gave higher edible yield than the unwrapped. Figure 5 shows the condition of lettuce stored 5 weeks after picking.

Results of experiment No. 2 (Table 2) indicate that lettuce stored for 17 days at 40°F and followed by storage for 22 days under Oxytrol CA showed higher mean scores for overall quality and gave significantly higher edible yield than that stored under normal atmosphere for 31 days. After 39 days, lettuce stored under normal atmosphere was a total loss, whereas 35.8 percent edible yield was obtained from lettuce stored under Oxytrol C.A.

* Although the O₂ control system was set at 4 and 5 percent, the chemical analysis indicated a range of 3-5.8%. Temperatures fluctuated between 34° and 36°F, although the control was set at 34°F.

Results of experiment No. 3 (Table 3) indicate that lettuce stored for 14 days under Oxytrol C.A. followed by 20 days at 40°F. under normal atmosphere showed higher mean scores for overall quality and gave significantly higher edible yield than the ones stored at similar temperatures under normal atmosphere.

Results of experiment No. 4 (Table 4) indicate that after six weeks of storage at 34-36°F. under normal atmosphere the lettuce was completely inedible, whereas 12.5 percent of edible yield was obtained from lettuce stored under Oxytrol C.A.

When the Oxytrol system was unopened for 4 weeks, the O₂ level dropped to 3 percent. This did not constitute an unaerobic condition.

Keeping time of up to 7 weeks under Oxytrol C.A. system is sufficient to meet anticipated overseas shipping time requirements.

REFERENCES

1. Lipton, W. J. Anatomical observations on russet spotting and pink rib of lettuce. Am. Soc. for Hort. Sci - Vol 78, 367-74, 1961.
2. Parsons, C. S. Extending the storage life of cabbage, celery, lettuce, and tomatoes aboard a Navy Supply Ship. USDA Marketing research Report No. 336, 1959.
3. Parsons, C. S., L.P. McCulloch, and R. C. Wright. Cabbage, celery, lettuce and tomatoes. Laboratory tests of storage methods, USDA Marketing Research Report No. 402, 1960.
4. Lipton, W. J. and Barger, W. R. Market quality of head lettuce in relation to delays between harvest and precooling and temperature after cooling. U.S. Dept. of Agr. Agr. Research Service. Hyattsville, Maryland. ARS 51-5, 1965.
5. Pratt, H. K., Morris, L. L., and Tucker, C. L. Temperature and lettuce deterioration. Proc. Comb. on Transport of Perishables. Univ. of Calif. Davis, 77-83, 1954.
6. Stewart, J. K. and Harvey, J. M. Temperatures, Relative Humidity, and Atmosphere composition in a mechanically refrigerated car and a trailer loaded with lettuce. ARS 51-13 U.S. Dept. of Agr. Agr. Res. Service. Fresno, California, 1967.
7. Bratley, C.O. Keeping quality of head lettuce as affected by holding temperature and various wrapping materials. U.S. Dept. Agr. Bur. Plant Ind., Soils, and Agr. Eng. H.T. and S. Office Rept. 151, 1945.
8. Rood, P. Relation of ethylene and post harvest temperature to brown spot of lettuce, Proc. Amer. Soc. Hort. Sci. 68: 296-303, 1956.
9. Watada, A. E., L. L. Morris, and L. Rappaport, Modified atmosphere effects on lettuce. Fruit and Veg. Perishables Handling Conf. Proc., Univ. Calif., 23-25: pp 82-85, 1964.
10. Parsons, C. S., J. E., Gates and D. H. Spalding, Quality of some fruits and vegetables after holding in nitrogen atmospheres. Amer. Soc. Hort. Sci. Proc. 84: 549-556, 1964.
11. Littlefield, N. A., B. N., Wanker, D. K. Salunkhe, and J. N. McGill Fungistatic effects of controlled atmospheres. Applied Microbiology, Vol. 14: 579-81, 1966.
12. Lipton, W. J. Market quality and rate of respiration of head lettuce held in low-oxygen atmospheres. Marketing research report No. 777. Agr. Res. Service U.S. Dept. of Agriculture, Washington, D. C., 1967.

13. Lugg, J. R. The Transfresh system of atmospheres for product transportation. Horticulture report No. 9, Dept. of Hort. Michigan State University, East Lansing, Michigan, p 58-61, 1969.
14. White, T. G. Summary of commodity holding study. Report No. 11. Oxytrol-A division of Occidental Petroleum Corporation, Burlingame, California, 1968.
15. Stewart, J. K. Effect of transit temperatures and modified atmospheres on market quality of lettuce shipped in nitrogen refrigerated and mechanically refrigerated trailers. Marketing research report No. 832, Agr. Res. Service, United States Dept. of Agr. Washington, D. C., 1968.
16. El-Mansy, H. I., D. K., Salunkhe, R. L. Hurst, and D. R. Walker, Effects of Pre- and post- harvest applications of 6 - purpurylaminopurine and of N6-Benzyladenine of physiological and chemical changes in lettuce. Horticultural Research, Vol. 7 No. 22 pp. 81-89, 1967.
17. Gorfien, H., A. R., Rahman, K. R. Johnson, and E. E. Anderson, Effect of a controlled atmosphere system on the storage life of lettuce. Technical report 70-23-FL, Food Laboratory FL-99, U.S. Army Natick Laboratories, Natick, Massachusetts, 1969.




Figure 1.

FRONT VIEW - OXYTROL MODEL O50 PORTABLE CONTAINER - OPERATING CONTROLS

LETTUCE EVALUATION
TEST # _____

SCORING CRITERIA	O.A. (W.W.L) = Overall Appearance - With Wrapper Leaves O.A. (Trimmed) = Overall Appearance - Trimmed																		Date _____																																											
	<table style="width: 100%; border: none;"> <tr> <td style="text-align: center;">* Field Fresh 9</td> <td style="text-align: center;">8</td> <td style="text-align: center;">Good-Minor Defects 7</td> <td style="text-align: center;">6</td> <td style="text-align: center;">Fair-Removable Defects 5</td> <td style="text-align: center;">4</td> <td style="text-align: center;">Poor-Generally Unsalable 3</td> <td style="text-align: center;">2</td> <td style="text-align: center;">In- edible 1</td> </tr> </table>																		* Field Fresh 9	8	Good-Minor Defects 7	6	Fair-Removable Defects 5	4	Poor-Generally Unsalable 3	2	In- edible 1	Duration _____																																		
	* Field Fresh 9	8	Good-Minor Defects 7	6	Fair-Removable Defects 5	4	Poor-Generally Unsalable 3	2	In- edible 1																																																					
	Decay - Pink Rib - Russet Spotting																		Treatment _____																																											
<table style="width: 100%; border: none;"> <tr> <td style="text-align: center;">Defects Absent 9</td> <td style="text-align: center;">8</td> <td style="text-align: center;">Trace 7</td> <td style="text-align: center;">6</td> <td style="text-align: center;">Slight 5</td> <td style="text-align: center;">4</td> <td style="text-align: center;">Moderate 3</td> <td style="text-align: center;">2</td> <td style="text-align: center;">Severe 1</td> </tr> </table>																		Defects Absent 9	8	Trace 7	6	Slight 5	4	Moderate 3	2	Severe 1	Carton # _____																																			
Defects Absent 9	8	Trace 7	6	Slight 5	4	Moderate 3	2	Severe 1																																																						
<table style="width: 100%; border: none;"> <tr> <td colspan="12" style="text-align: center;">Top Layer</td> <td colspan="8" style="text-align: center;">Bottom Layer</td> </tr> <tr> <td style="text-align: center;">1</td><td style="text-align: center;">2</td><td style="text-align: center;">3</td><td style="text-align: center;">4</td><td style="text-align: center;">5</td><td style="text-align: center;">6</td><td style="text-align: center;">7</td><td style="text-align: center;">8</td><td style="text-align: center;">9</td><td style="text-align: center;">10</td><td style="text-align: center;">11</td><td style="text-align: center;">12</td> <td style="text-align: center;">13</td><td style="text-align: center;">14</td><td style="text-align: center;">15</td><td style="text-align: center;">16</td><td style="text-align: center;">17</td><td style="text-align: center;">18</td><td style="text-align: center;">19</td><td style="text-align: center;">20</td><td style="text-align: center;">21</td><td style="text-align: center;">22</td><td style="text-align: center;">23</td><td style="text-align: center;">24</td> </tr> </table>																		Top Layer												Bottom Layer								1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	Chamber # _____
Top Layer												Bottom Layer																																																		
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24																																							
DEFECTS																		Name _____																																												
O.A. (W.W.L)																																																														
O.A. (Trimmed)																																																														
Decay (Slime, mold, etc.)																																																														
Pink Rib																																																														
Russet Spotting																																																														

* The following are considered defects: Discoloration such as browning, darkening, black, pink, red spotting or the presence of any color not typical of the fresh product; physical damage such as bruises and skin breakage; diseases such as mold, slime and soft rot; wrinkled surface; wilt; tip burn; spotting; dehydration and any other sign not typical of the fresh product.

Differences in appearance of butt color _____

Other Comments _____

Figure 2. Technological Evaluation Sheet

Figure 3. Effect of Storage Conditions on the Quality of Lettuce



Unwrapped lettuce stored under Oxytrol Controlled Atmosphere System 5 Weeks after Picking(1 week transit & 4 weeks at NLABS).



Unwrapped lettuce stored under Normal atmosphere at approximately 34°F., 5 weeks after picking(1 week transit & 4 weeks at NLABS).



Wrappped lettuce stored under Oxytrol Controlled Atmosphere System, 5 weeks after picking (1 week transit & 4 weeks at NLABS).



Wrappped lettuce stored under normal atmosphere at approximately 34°F., 5 weeks after picking(1 week transit & 4 weeks at NLABS).



Bottom layer of a 4 fiberboard boxes stack of lettuce stored under normal atmosphere conditions at approximately 34°F., 5 weeks after picking (1 week transit & 4 weeks at NLABS).

Table 1. Effect of Storage Conditions on the Quality of Lettuce Stored 5 to 7 Weeks After Picking

Storage Conditions	Storage Time After Picking Weeks	Percent Edible Yield		Mean Technological Scores			
		Wrapped	Unwrapped	Defects		Overall Appearance	
				Wrapped	Unwrapped	Wrapped	Unwrapped
Oxytrol 4.4% O ₂ at 34°F.	5	66.0*	50.5*	7.0	6.6	8.3	8.0
Normal atmosphere at 34°F.	5	14.2*	12.0*	3.0	2.2	1.3	2.3
Oxytrol 4.5% O ₂ at 34°F. for 5 Weeks plus 1 week under normal atmosphere at 40°F.	6	60.6*	28.8*	5.9	4.2	6.0	5.8
Normal atmosphere at 40°F.	6	0*	0*	1	1	1	1
Oxytrol 4.4% O ₂ at 34°F.	7	27.5*	27.0*	6.1	6.0	7.1	7.2
Normal atmosphere at 34°F.	7	0*	0*	1	1	1	1
Oxytrol 4.5% O ₂ at 34°F. for 5 weeks plus 2 weeks under normal atmosphere at 40°F	7	21.4	16.0	3.9	3.8	4.0	3.9

* Significant at the 1 percent level

Table 2. Effect of Storage Conditions on the Quality of Lettuce Stored for 21 to 39 Days After Purchase

Storage Conditions	Storage Time after purchase	*Percent Edible Yield	Mean Technological Scores	
			Overall Quality	Defects
	Days			
Normal atmosphere at 40°F. for 17 days plus 4 days, Oxytrol 5-8% O ₂ at 34°F.	21	60.8*	7.2	7.5
Normal atmosphere at 40°F. plus 4 days at 36°F.	21	55.6*	6.1	6.0
Normal atmosphere at 40°F. for 17 days plus 14 days Oxytrol 5.8% O ₂ at 34°F.	31	43.5*	5.3	5.5
Normal atmosphere at 40°F. for 17 days plus 14 days at 34°F.	31	25.3*	4.3	4.1
Normal atmosphere at 40°F. for 17 days plus 22 days Oxytrol 5-5% O ₂ at 34°F.	39	35.8*	4.3	4.5
Normal atmosphere at 40°F. for 17 days plus 22 days at 34°F.	39	0	1	1

* Significant at the 1 percent level.

Table 3. Effect of Storage Conditions on the Quality of Lettuce Stored for 14 to 34 days after Purchase

Storage Conditions	Storage Time after purchase	Percent Edible Yield	Mean Technological Scores	
			Overall Quality	Defects
	Days			
Oxytrol 5±7% O ₂ at 34°F.	14	57.4*	8.0	8.3
Normal atmosphere at 34°F.	14	44.4*	7.0	7.4
Oxytrol 5±7% O ₂ at 34°F. plus 8 days at 40°F. under normal atmosphere.	22	55.8*	5.3	6.0
Normal atmosphere at 34°F. for 14 days plus 8 days at 40°F.	22	21.0*	3.4	4.0
Oxytrol 5±7% O ₂ at 34°F. plus 20 days at 40°F. under normal atmosphere.	34	40.2*	4.9	5.1
Normal atmosphere at 34°F. for 14 days plus 20 days at 40°F.	34	10.5*	3.4	3.7

* Significant at the 1 percent level

Table 4. Effect of Storage Conditions on the Quality of Lettuce Stored for 2 to 6 Weeks after Purchase

Storage Conditions	Storage Time after Purchase Weeks	Percent Edible Yield	Mean Technological Scores	
			Overall Quality	Defects
Oxytrol 5-7% O ₂ at 34°F.	2	57.4*	7.4	7.3
Normal atmosphere at 34°F.	2	46.5*	7.0	6.8
Oxytrol 3-4% O ₂ at 34°F.	6	12.5*	2.5	2.9
Normal atmosphere at 34°F	6	0 *	1	1

* Significant at the 1 percent level.

FOOD LABORATORY DISTRIBUTION LIST - PLANT PRODUCTS

Copies

- 1 - Commanding General
US Army Medical Research and
Development Command
ATTN: MEDDH-SI
Washington, D. C. 20315

- 2 - Commanding General
US Army Test and Evaluation
Command
ATTN: AMSTE-BC
Aberdeen Proving Ground,
Maryland 21005

- 1 - Commanding General
US Army Combat Development
Command
Combat Service Support Group
Fort Lee, Virginia 23801

- 1 - Commanding General
US Army Combat Development
Command
ATTN: GDCMR-O
Fort Belvoir, Virginia 22060

- 1 - Commanding General
US Army Materiel Command
ATTN: AMCRD-JI
Department of the Army
Washington, D. C. 20315

- 2 - Commanding Officer
Edgewood Arsenal
ATTN: SMUEA-TSTI-TL
Edgewood Arsenal, Maryland
21010

- 1 - Commanding Officer
US Army Combat Development
Command
Supply Agency
ATTN: GDCSA-R
Fort Lee, Virginia 23801

Copies

- 1 - Commanding Officer
US Army Medical Nutrition
Laboratory
Fitzsimons General Hospital
Denver, Colorado 80240

- 1 - Commander
Defense Personnel Support
Center
ATTN: Directorate of
Subsistence, DPSC-STC
2800 South 20th Street
Philadelphia, Pennsylvania
19101

- 1 - Commandant of the Marine Corps
Code AO4D
Washington, D. C. 20380

- 1 - Commandant of the Marine Corps
Code CDE
Washington, D. C. 20380

- 2 - Executive Secretary
Interdepartmental Committee on
Radiation Preservation of
Food
Consumer Products Division 623
Business and Defense Service
Administration
US Department of Commerce
Washington, D. C. 20230

- 2 - Director
Development Center
Marine Corps Development and
Education Command
ATTN: Combat Service Support
Division
Quantico, Virginia 22134

Copies

- 1 - Director
Division of Biology and
Medicine
US Atomic Energy Commission
Washington, D. C. 20545
- 1 - Director
US Army Advanced Materiel
Concepts Agency
Washington, D. C. 20315
- 1 - Chief, Life Sciences Division
Army Research Office
Office of Chief of Research
and Development
Washington, D. C. 20310
- 3 - Office of the Coordinator of
Research
University of Rhode Island
Kingston, Rhode Island 02881
- 1 - Dr. Herbert E. Hall
Chief, Food Microbiology
National Center for Urban and
Industrial Health
Food Protection Research
222 East Central Parkway
Cincinnati, Ohio 45202
- 1 - Library Southern Utilization
Research and Development
Division
Agricultural Research Service
US Department of Agriculture
PO Box 19687
New Orleans, Louisiana 70119

Copies

- 1 - Stimson Library
ATTN: Documents Librarian
US Army Medical Field Service
School
Brooke Army Medical Center
Fort Sam Houston, Texas 78234
- 2 - Quartermaster School Library
US Army Quartermaster School
Fort Lee, Virginia 23801
- 2 - Technical Library
USACDC Institute of Land
Combat
301 Taylor Drive
Alexandria, Virginia 22314
- 1 - US Department of Agriculture
Division of Acquisitions
National Agriculture Library
Washington, D. C. 20250
- 2 - US Army Research Office
ATTN: Technical Library
3045 Columbia Pike
Arlington, Virginia 22204
- 2 - Headquarters 12th Support
Brigade
ACofS Services
ATTN: Food Advisor
Fort Bragg, North Carolina
28307
- 4 - Exchange and Gift Division
Library of Congress
Washington, D. C. 20540

Copies

- 2 - Commanding General
Defense Personnel Support
Center
ATTN: Jim Burnham
2800 South 20th Street
Philadelphia, PA. 19101
- 1 - Commanding General
Defense Personnel Support
Center
ATTN: Mr. Arthur Solomon
2800 South 20th Street
Philadelphia, PA. 19101
- 2 - US Army Natick Laboratories
Office
Defense Personnel Support Center
ATTN: Officer in Charge
2800 South 20th Street
Philadelphia, PA. 19101
- 1 - Commander
Naval Supply Systems Command
(SUP 063), Navy Department
Washington, D.C. 20360
- 1 - Commanding Officer
US Navy Subsistence Office
Washington Navy Yard
Washington, D.C. 20390
- 1 - Commander
Naval Ships Engineering Center
ATTN: Mr. Arthur S. Childers
Code 6031/32
Washington, D.C. 20360
- 1 - Chief of Naval Research (466)
Navy Department
Washington, D.C. 20360
- 1 - Commandant of the Marine Corps
Headquarters US Marine Corps
ATTN: Research & Development Section
Washington, D.C. 20360
- 5 - Oxytrol Corporation
ATTN: Dr. Thomas White
895 Milton Road
P.O. Box 1488 Station A
Burlingame, California 94010

Copies

- 1 - Director
Defense Supply Agency (DSAH-N)
Cameron Station
Alexandria, Virginia 22300
- 1 - Chief, US Army Food Service
Center
1819 West Pershing Road
Chicago, Illinois 60609
- 1 - Commanding General
Headquarters, US Army, Alaska
APO Seattle 98749
- 1 - Director of Research and
Development, Headquarters
US Air Force
Washington, D.C. 20360
- 1 - Market Quality Research
Division
Agricultural Research
Service, USDA
Belle Mead, New Jersey 08502
- 1 - Market Quality Research
Division
Horticultural Crops Research
Branch
Agricultural Research Service,
Plant Industry Station
Beltsville, Maryland 20705
- 1 - Commander
Defense Supply Agency
Subsistence Regional
Headquarters, DPSC
ATTN: Chief, Fruits and
Vegetable Branch
312 N. Spring Street
Los Angeles, California 90012
- 2 - Directorate for Food Service
Office of the Assistant Secretary
of Defense
ATTN: Captain James A. Warren
SC USN
Room 2B 323 - The Pentagon
Washington, D.C. 20301

Copies

- 2 - Dr. Frank R. Fisher, AMPS
National Academy of Sciences
National Research Council
2101 Constitution Avenue
Washington, D. C. 20418
- 1 - Dr. Irving J. Pflug
Mayo Memorial Hospital
University of Minnesota
Minneapolis, Minnesota 56524
- 1 - Dr. Joseph R. Wagner
Processing Technology Investigations
Vegetable Laboratory
Western Utilization Research and
Development Division
US Department of Agriculture
Albany, California 94706
- 1 - Commander
Headquarters, US Army Materiel
Command
Chief, Ind. & General Equipment Office
Development Directorate
Washington, D. C. 20315
- 1 - Commander, Military Traffic Manage-
ment and Terminal Service
ATTN: MTMTS-F
Washington, D. C. 20315
- 1 - Transfresh Corporation
32 E. Alisal Street
ATTN: Mr. J. Lamp
Salinas, California 43901
- 1 - Commanding General
Headquarters, US Army Materiel
Command
Washington, D.C. 20315

Copies

- 1 - U.S.D.A.
Agricultural Research Service
Transportation and Facilities
Research Division
Federal Center Building
ATTN: Mr. R. E. Guilfooy
Hyattsville, Maryland 20782
- 1 - U.S.D.A.
Transportation & Facilities
Research Division
Agricultural Research Service
Federal Center Building
ATTN: Mr. D. R. Stokes
Hyattsville, Maryland 20782
- 1 - Kaiserslautern Cold Stores
US Army Supply and
Maintenance Agency
ATTN: Major Gagliardone
APO 09054 New York, N.Y.
- 1 - The Commander in Chief
Headquarters, US Army Europe
and Seventh Army
APO New York 09403
- 1 - US Army, S&MA Zweibrücken
ATTN: Mr. R. D. Gentzler
APO New York, N.Y.
- 1 - Kaiserslautern Cold Stores
US Army Supply and
Maintenance Agency
ATTN: Mr. Sidney Novitch
APO 09054 New York, N.Y.
- 2 - Commanding General
US Army Medical Research and
Development Command
Main Navy Building
Washington, D. C. 20315
- 1 - Office of Technical Services
U. S. Department of Commerce
Washington, D. C. 20360

Copies

- 1 - Commander
US Army Biological Laboratories
ATTN: Technical Library
Fort Detrick
Frederick, Maryland 21701
- 1 - US Army Research Office
3045 Columbia Pike
Arlington, Virginia 22204
- 1 - Defense Supply Agency, DSA-WT
ATTN: Mr. Roland Priesman
Cameron Station, Virginia 22314
- 1 - Dr. John M. Harvey
US Department of Agricultural
Agricultural Research Service
Market Quality Research Division
2021 South Peach Avenue
Fresno, California 93727
- 2 - DSA Subsistence Regional Hdqs,
Oakland
ATTN: Mr. William S. Chu
Deputy Chief Supp.
Alameda, California 94501
- 1 - National Bureau of Standards
Building Research Division
ATTN: Mr. Clinton Philips
Room B-104, Building 226
Washington, D. C. 20234
- 1 - Commanding Officer
U. S. Naval Supply Depot Yokosuka
ATTN: Lt R. E. Bill
Fleet Post Office, Seattle 98762

Copies

- 1 - Commanding Officer
U. S. Naval Supply Depot Yokosuka
ATTN: Mr. Kelly Woolums
Fleet Post Office, Seattle 98762
- 1 - Naval Logistics Engineering Group
ATTN: Mr. J. Brugh
Cheatham Annex
Williamsburg, Virginia 23185
- 1 - US Army Veterinary Det, Japan
Camp Zama
ATTN: Colonel J. Grogan
APO 96343, San Francisco
- 1 - Commanding Officer
Naval Supply Center
ATTN: Commander W. C. Hamilton
Norfolk, Virginia 23512
- 1 - Research Development and
Engineering Directorate
Science Technology Division
ATTN: Dr. H. M. El-Bisi
Nassif Bldg - Rm 232
Washington, D. C. 20315

FOOD LABORATORY INTERNAL DISTRIBUTION LIST- PLANT PRODUCTS

Copies

- 25 - Chief, Technical Plans Office, NLABS
(20 for transmittal to Defense Documentation Center)
- 2 - Technical Library, NLABS
- 10 - Program Coordination Office, Food Laboratory, NLABS
- 7 - Division Chiefs, Food Laboratory, NLABS
- 2 - Marine Liaison Officer, NLABS
- 3 - Air Force Liaison Officer, NLABS
- 1 - Director, Earth Sciences Laboratory, NLABS
- 2 - Director, General Equipment and Packaging Laboratory, NLABS
- 3 - Director, Pioneering Research Laboratory, NLABS
- Project Officer and Alternate Project Officer, Food Laboratory, NLABS
- 60 - Food Laboratory, NLABS

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)

US Army Natick Laboratories
Natick, Massachusetts 01760

2a. REPORT SECURITY CLASSIFICATION

Unclassified

2b. GROUP

3. REPORT TITLE

Storage Life of Lettuce as Affected by Controlled Atmosphere System

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

5. AUTHOR(S) (First name, middle initial, last name)

Abdul R. Rahman, Glenn Schafer, George R. Taylor, Donald E. Westcott

6. REPORT DATE

January 1970

7a. TOTAL NO. OF PAGES

20

7b. NO. OF REFS

17

8a. CONTRACT OR GRANT NO.

b. PROJECT NO.

2270.3

c.

d.

9a. ORIGINATOR'S REPORT NUMBER(S)

70-48-FL

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

FL-106

10. DISTRIBUTION STATEMENT

This document has been approved for public release and sale; its distribution is unlimited.

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

US Army Natick Laboratories
Natick, Massachusetts 01760

13. ABSTRACT

The effect of Oxytrol controlled atmosphere system on the shelf life of lettuce was evaluated. Lettuce stored under Oxytrol at oxygen levels ranging from 3 to 5.8 percent and temperatures from 34 to 36°F for 2 to 7 weeks gave significantly higher edible yield and showed higher mean scores for overall quality than lettuce stored under normal atmosphere at similar temperatures.

DD FORM 1473
1 NOV 66

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS
OBSOLETE FOR ARMY USE.

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Controlled atmosphere	6		5,10			
Oxytrol system	6		5,10			
Lettuce	7		9			
Storage	7		8,4			
Protection	7		8			
Military bases	4		4			
Transportation	4		8			